


CLAIMS

1. A compound of formula (1), or a pharmaceutically acceptable salt, solvate or *in vivo*
 5 hydrolysable ester thereof:



- wherein Y is selected from a bond, -S-, -O-, -NR⁵-, -CF₂-CH₂-, -CF₂CF₂-, -CONR⁵-,
 10 phenyl or heteroaryl.
 wherein R¹ is a group selected from C₃₋₇carbocyclyl, C₁₋₈alkyl, C₂₋₆alkenyl and C₂₋₆alkynyl;
 wherein the group is optionally substituted by 1, 2 or 3 substituents independently selected
 from fluoro, nitrile, -OR⁴, -NR⁵R⁶, -CONR⁵R⁶, -COOR⁷, -NR⁸COR⁹, -SR¹⁰, -SO₂R¹⁰,
 -SO₂NR⁵R⁶, -NR⁸SO₂R⁹, phenyl or heteroaryl; wherein phenyl and heteroaryl are optionally
 15 substituted by 1, 2 or 3 substituents independently selected from halo, cyano, nitro, -OR⁴, -
 NR⁵R⁶, -CONR⁵R⁶, -COOR⁷, -NR⁸COR⁹, -SR¹⁰, -SO₂R¹⁰, -SO₂NR⁵R⁶, -NR⁸SO₂R⁹,
 C₁₋₆alkyl and trifluoromethyl;
 wherein R² is C₃₋₇carbocyclyl, optionally substituted by 1, 2 or 3 substituents independently
 selected from fluoro, -OR⁴, -NR⁵R⁶, -CONR⁵R⁶, -COOR⁷, -NR⁸COR⁹, -SR¹⁰, -SO₂R¹⁰,
 20 -SO₂NR⁵R⁶, -NR⁸SO₂R⁹;
 or R² is a 3-8 membered ring optionally containing 1, 2 or 3 atoms selected from O, S, -NR⁸
 and whereby the ring is optionally substituted by C₁₋₃alkyl or fluoro;
 or R² is a phenyl or heteroaryl, each of which is optionally substituted by 1, 2 or 3
 substituents independently selected from halo, cyano, nitro, -OR⁴, -NR⁵R⁶, -CONR⁵R⁶, -
 25 NR⁸COR⁹, -SO₂NR⁵R⁶, -NR⁸SO₂R⁹, C₁₋₆alkyl and trifluoromethyl;
 or R² is a group selected from C₁₋₈alkyl, C₂₋₆alkenyl or C₂₋₆alkynyl wherein the group is
 substituted by 1, 2 or 3 substituents independently selected from hydroxy, amino, C₁₋₆alkoxy,
 C₁₋₆alkylamino, di(C₁₋₆alkyl)amino, N-(C₁₋₆alkyl)-N-(phenyl)amino, N-C₁₋₆alkylcarbonyl,

-33-

N,N-di(C₁₋₆alkyl)carbamoyl, *N*-(C₁₋₆alkyl)-*N*-(phenyl)carbamoyl, carboxy, phenoxycarbonyl, -NR⁸COR⁹, -SO₂R¹⁰, -SO₂NR⁵R⁶ and -NR⁸SO₂R⁹;

wherein R³ is hydrogen or independently R²;

R⁴ is hydrogen or a group selected from C₁₋₆alkyl and phenyl, wherein the group is optionally substituted by 1 or 2 substituents independently selected from halo, phenyl, -OR¹¹ and -NR¹²R¹³;

R⁵ and R⁶ are independently hydrogen or a group selected from C₁₋₆alkyl and phenyl wherein the group is optionally substituted by 1, 2 or 3 substituents independently selected from halo, phenyl, -OR¹⁴, -NR¹⁵R¹⁶, -COOR¹⁴, -CONR¹⁵R¹⁶, -NR¹⁵COR¹⁶, -SO₂R¹⁰, -SONR¹⁵R¹⁶ and NR¹⁵SO₂R¹⁶

or

R⁵ and R⁶ together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocyclic ring system optionally containing a further heteroatom selected from oxygen and nitrogen atoms, which ring is optionally substituted by 1, 2 or 3 substituents independently selected from phenyl, -OR¹⁴, -COOR¹⁴, -NR¹⁵R¹⁶, -CONR¹⁵R¹⁶, -NR¹⁵COR¹⁶, -SO₂R¹⁰, -SONR¹⁵R¹⁶, NR¹⁵SO₂R¹⁶ or C₁₋₆alkyl (optionally substituted by 1 or 2 substituents independently selected from halo, -NR¹⁵R¹⁶ and -OR¹⁷ groups);

R¹⁰ is hydrogen or a group selected from C₁₋₆alkyl or phenyl, wherein the group is optionally substituted by 1, 2 or 3 substituents independently selected from halo, phenyl, -OR¹⁷ and -NR¹⁵R¹⁶; and each of R⁷, R⁸, R⁹, R¹¹, R¹², R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷ is independently hydrogen, C₁₋₆alkyl or phenyl;

R^x is trifluoromethyl, -NR⁵R⁶, phenyl, naphthyl, monocyclic or bicyclic heteroaryl wherein a heteroring may be partially or fully saturated and one or more ring carbon atoms may form a carbonyl group, and wherein each phenyl or heteroaryl group is optionally substituted by 1, 2 or 3 substituents independently selected from halo, cyano, nitro, -OR⁴, -NR⁵R⁶, -CONR⁵R⁶, -COR⁷, -COOR⁷, -NR⁸COR⁹, -SR¹⁰, -SO₂R¹⁰, -SO₂NR⁵R⁶, -NR⁸SO₂R⁹, C₁₋₆alkyl or trifluoromethyl;

or R^x is a group selected from C₃₋₇carbocyclyl, C₁₋₈alkyl, C₂₋₆alkenyl and C₂₋₆alkynyl whereby the group is optionally substituted by 1, 2 or 3 substituents independently selected from halo, -OR⁴, -NR⁵R⁶, -CONR⁵R⁶, -COR⁷, -COOR⁷, -NR⁸COR⁹, -SR¹⁰, -SO₂R¹⁰, -SO₂NR⁵R⁶, -NR⁸SO₂R⁹, phenyl or heteroaryl; and wherein each phenyl or heteroaryl group is optionally substituted by 1, 2 or 3 substituents independently selected from halo, cyano, nitro, -OR⁴, -

-34-

NR⁵R⁶, -CONR⁵R⁶, -COR⁷, -COOR⁷, -NR⁸COR⁹, -SR¹⁰, -SO₂R¹⁰, -SO₂NR⁵R⁶,
-NR⁸SO₂R⁹, C₁₋₆alkyl or trifluoromethyl;

2. A compound, or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof according to claim 1 wherein R² is C₁₋₈alkyl optionally substituted by 1 or 2 hydroxy substituents.
3. A compound, pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof according to claim 1 wherein R¹ is benzyl or -CH₂CH₂OPh, or CH₂CH₂Ph wherein in each case the phenyl ring is optionally substituted by 1, 2 or 3 substituents independently selected from fluoro, chloro, bromo, methoxy, methyl and trifluoromethyl.
4. A compound, pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof wherein R³ is hydrogen.
5. A compound, pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof wherein Y is selected from a bond, -S-, and -CF₂-CH₂- and -CH₂-CH₂-.
6. A compound, pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof wherein R^x is methyl, 1-methylimidazolyl, 1,2-dimethylimidazolyl, *N,N*-dimethylamino, azetidiny, pyrrolidiny, morpholinyl, piperidinyl and trifluoromethyl
7. A compound selected from the group consisting of:
N-[4-[(2,3-difluorophenyl)methyl]thio]-6-[(1*R*)-2-hydroxy-1-methylethyl]amino]-1,3,5-triazin-2-yl]-methanesulfonamide; and
N-[4-[(2,3-difluorophenyl)methyl]thio]-6-[(1*R*)-2-hydroxy-1-methylethyl]amino]-1,3,5-triazin-2-yl]-1-azetidinesulfonamide, *N*-[4-[(2,3-difluorophenyl)methyl]thio]-6-[(1*R*)-2-hydroxy-1-methylethyl]amino]-1,3,5-triazin-2-yl]-methanesulfonamide
N-[4-[(2,3-difluorophenyl)methyl]thio]-6-[(1*R*)-2-hydroxy-1-methylethyl]amino]-1,3,5-triazin-2-yl]-1-azetidinesulfonamide
4-morpholinesulfonamide, *N*-[4-[(2,3-difluorophenyl)methyl]thio]-6-[(1*R*)-2-hydroxy-1-methylethyl]amino]-1,3,5-triazin-2-yl]-

-35-

methanesulfonamide, *N*-[4-[[2-(2,3-difluorophenoxy)ethyl]thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]-1,3,5-triazin-2-yl]-

methanesulfonamide, 1,1,1-trifluoro-*N*-[4-[[*(1R)*-2-hydroxy-1-methylethyl]amino]-6-(2-phenylethyl)-1,3,5-triazin-2-yl]- or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof.

8. A compound, or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof according to any one of claims 1 to 7 for use as a medicament.

10 9. A compound, or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof according to any one of claims 1 to 7 for use as a medicament for the treatment of asthma, allergic rhinitis, COPD, inflammatory bowel disease, osteoarthritis, osteoporosis, rheumatoid arthritis, or psoriasis..

15 10. A compound, or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof according to any one of claims 1-7, for use as a medicament for the treatment of cancer.

11. The use of a compound, or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof, according to any one of claims 1 to 7 in the manufacture of a medicament for the treatment of human diseases or conditions in which modulation of chemokine receptor activity is beneficial.


12. The use of a compound, or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof, according to any one of claims 1 to 7 in the manufacture of a medicament for the treatment of asthma, allergic rhinitis, COPD, inflammatory bowel disease, irritable bowel syndrome, osteoarthritis, osteoporosis, rheumatoid arthritis, or psoriasis.

30 13. The use of a compound, or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof, according to any one of claims 1 to 7 in the manufacture of a medicament for the treatment of cancer.

-36-

14. A pharmaceutical composition comprising a compound, or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof according to any one of claims 1 to 7; and a pharmaceutically-acceptable diluent or carrier.

5 15. A process for the preparation of a compound according to claim 1 or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof, which comprises the steps of:
treating a compound of formula (2):



wherein Y, R¹, R² and R³ are as defined in formula (1) with a sulfonamide of formula R^xSO₂NH₂ where R^x is as defined in formula (1);

15 and optionally thereafter, one or more of steps (i), (ii), (iii), (iv), or (v) in any order:

i) removing any protecting groups;

ii) converting the compound of formula (1) into a further compound of formula (1)

iii) forming a salt

iv) forming a prodrug

20 v) forming an *in vivo* hydrolysable ester.

16. A combination therapy which comprises administering a compound of formula (1) or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof, or a pharmaceutical composition or formulation comprising a compound of formula (1),
25 concurrently or sequentially with other therapy and/or another pharmaceutical agent.

17. A combination therapy as claimed in claim 16 for the treatment of asthma, allergic rhinitis, COPD, inflammatory bowel disease, irritable bowel syndrome, osteoarthritis, osteoporosis, rheumatoid arthritis, or psoriasis.

-37-

18. A combination therapy as claimed in claim 16 for the treatment of cancer.

19. A pharmaceutical composition which comprises a compound of formula (1) or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof, in conjunction
5 with another pharmaceutical agent.

20. A pharmaceutical composition as claimed in claim 19 for the treatment of asthma, allergic rhinitis, COPD, inflammatory bowel disease, irritable bowel syndrome, osteoarthritis, osteoporosis, rheumatoid arthritis, or psoriasis.

10

21. A pharmaceutical composition as claimed in claim 19 for the treatment of cancer.